

Total No. of Printed Pages—8

5 SEM TDC CHMH (CBCS) C 12

2 0 2 1

(Held in January/February, 2022)

CHEMISTRY

(Core)

Paper : C-12

(Physical Chemistry)

Full Marks : 53

Pass Marks : 21

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

1. Choose the correct answer from the following : 1×4=4

(a) The degeneracy of rotational level of a diatomic molecule having energy $\frac{h^2}{4\pi^2 I}$ is

(i) 0

(ii) 1

(iii) 2

(iv) 3

- (b) Vibrational transition exists in
- (i) infrared region
 - (ii) microwave region
 - (iii) visible region
 - (iv) radio-frequency region
- (c) The degeneracy of a particle of mass m confined in a three-dimensional box having energy level equal to $\frac{14h^2}{8ma^2}$ is
- (i) 7
 - (ii) 14
 - (iii) 6
 - (iv) 8
- (d) In photosynthesis, chlorophyll acts as a
- (i) catalyst
 - (ii) photosensitizer
 - (iii) photoinhibitor
 - (iv) All of the above

2. Answer any *four* questions from the following : 2×4=8

- (a) Microwave studies are done only in gaseous state. Explain.

(3)

- (b) Explain why the nuclei H^1 and ^{13}C are suitable for NMR investigation.
- (c) Write a short note on fingerprint region.
- (d) What is chemiluminescence? Give one example.
- (e) Show that the functions $\psi_1 = \left(\frac{1}{2\pi}\right)^{1/2}$ and $\psi_2 = \left(\frac{1}{\pi}\right)^{1/2} \cos x$ in the interval $x = 0$ to $x = 2\pi$ are orthogonal to each other.
- (f) Show that $\sin 4x$ is an eigenfunction of the operator $\frac{d^2}{dx^2}$. Find the eigenvalue.

UNIT—I

3. Answer any *four* questions from the following : 4×4=16

- (a) What are normalized and orthogonal wave functions? For the function $\psi(\theta) = \sin \theta$, where the variable θ changes continuously from 0 to 2π , determine whether it is normalized or not. If it is not, find the normalization factor.

1+2+1=4

- (b) ψ_i and ψ_j represent the wave function corresponding to two different states of a particle moving freely in a one-dimensional box. Show that they are orthogonal to each other.

4

- (c) Consider a particle of mass m confined in a two-dimensional box of edge lengths a and b . Find the energy and wave functions by solving the Schrödinger's equation. The potential energy

$$V(x, y) = 0, \text{ for } 0 \leq x \leq a \text{ and } 0 \leq y \leq b \\ = \infty, \text{ elsewhere}$$

Also write the expression for energy when $a = b$.

3+1=4

- (d) (i) What does the term 'degenerate levels' mean? Determine the degree of degeneracy of the level $\frac{17h^2}{8ma^2}$ of a particle in a cubical box.

1+1=2

- (ii) Form Schrödinger wave equation for a one-dimensional simple harmonic oscillator.

2

- (e) (i) The distance between the atoms of a diatomic molecule is r and its reduced mass is μ . If the angular momentum is L and moment of inertia is I , then prove that kinetic energy $T = \frac{L^2}{2\mu r^2}$. 3

- (ii) Write the expression for energy for a rigid rotator. 1

- (f) (i) Write down the Schrödinger wave equation in polar form for H-atom. 1½

- (ii) Calculate the most probable distance r_{mp} of the electron from the nucleus in the ground state of hydrogen atom, given that the normalized ground state wave function is

$$\psi_{1s} = \frac{1}{\sqrt{\pi}a_0^{3/2}} e^{(-r/a_0)}$$

Given $a_0 = 0.529 \text{ \AA}$. 2½

- (g) (i) Write down the equation showing Hamiltonian operator for one-dimensional harmonic oscillator. 2

(6)

- (ii) Sketch the variation of radial probability density against the distance from the nucleus for 2s state for hydrogen atom. 2

UNIT—II

4. Answer any two questions from the following : $8 \times 2 = 16$

- (a) (i) Show that the lines in the rotational spectrum of a diatomic molecule are equispaced under the rigid rotator approximation. 3

- (ii) A transition from $J=0$ to $J=1$ in the rotational spectrum of CO corresponds to 3.84235 cm^{-1} . Calculate the moment of inertia and bond length. $2+2=4$

- (iii) Write the selection rule for rotational spectra. 1

- (b) (i) Show that the frequency of the absorbed radiation in pure vibrational spectra is equal to the fundamental frequency of vibration ν_0 of the molecule. $2\frac{1}{2}$

- (ii) Prove that the ratio of wave numbers of fundamental, first overtone and second overtone is approximately 1:2:3. 2½
- (iii) Roughly sketch the fundamental modes of vibrations of CO₂ and show the infrared active vibrations. 3
- (c) (i) State and explain Franck-Condon principle. 3
- (ii) Explain why TMS is used as a reference substance in NMR spectroscopy. 2
- (iii) Calculate the NMR frequency (in MHz) of the proton (¹H) in a magnetic field of intensity 1.4092 tesla, given that $g_N = 5.585$ and $\mu_N = 5.05 \times 10^{-27} \text{ JT}^{-1}$. 2

Or

Briefly discuss Born-Oppenheimer approximation.

- (iv) Write any one difference between fluorescence and phosphorescence. 1

UNIT—III

5. Answer any *two* questions from the following : $4\frac{1}{2}\times 2=9$

(a) State and explain Lambert-Beer law. Write the significance of molar extinction coefficient. $4\frac{1}{2}$

(b) Explain the term 'quantum yield'. Discuss briefly the reasons for high and low quantum yields. $1\frac{1}{2}+3=4\frac{1}{2}$

(c) What is photochemical equilibrium? Give example of a photochemical equilibrium in which only one reaction is light sensitive. Deduce an expression for equilibrium constant of a photochemical equilibrium. $1+1+2\frac{1}{2}=4\frac{1}{2}$

★ ★ ★